

Treatment of Childhood and Adolescent Depression

MOLLY S. CLARK, PhD; KATE L. JANSEN, PhD; and J. ANTHONY CLOY, MD

University of Mississippi Medical Center, Jackson, Mississippi

Major depressive disorder in children and adolescents is a common condition that affects physical, emotional, and social development. Risk factors include a family history of depression, parental conflict, poor peer relationships, deficits in coping skills, and negative thinking. Diagnostic criteria are the same for children and adults, with the exception that children and adolescents may express irritability rather than sad or depressed mood, and weight loss may be viewed in terms of failure to reach appropriate weight milestones. Treatment must take into account the severity of depression, suicidality, developmental stage, and environmental and social factors. Cognitive behavior therapy and interpersonal therapy are recommended for patients with mild depression and are appropriate adjuvant treatments to medication in those with moderate to severe depression. Pharmacotherapy is recommended for patients with moderate or severe depression. Tricyclic antidepressants are not effective in children and adolescents. Antidepressants have a boxed warning for the increased risk of suicide; therefore, careful assessment, follow-up, safety planning, and patient and family education should be included when treatment is initiated. (*Am Fam Physician*. 2012;85(5):442-448. Copyright © 2012 American Academy of Family Physicians.)

► Patient information:

A handout on depression is available at <http://familydoctor.org/641.xml>.

The prevalence of depression is estimated to be 2.8 percent in children younger than 13 years and 5.6 percent in adolescents 13 to 18 years of age.¹ The incidence of depression among children and adolescents is of great concern because of the acute and lasting consequences associated with depressive disorders. Approximately 60 percent of adolescents with depression have recurrences throughout adulthood.² Furthermore, adults with a history of adolescent depression have a higher rate of suicide than those without such a history.² Adolescent-onset depression has been associated with abuse and neglect³; poor academic performance; substance use; early pregnancy; and disruptions in social, employment, and family settings into adulthood.⁴⁻⁶ Although the prevalence of adolescent depression is high, it is significantly underdiagnosed and undertreated.⁷ Because of the lack of mental health care professionals, family physicians are often responsible for detecting and treating childhood and adolescent depression.⁸⁻¹⁰

Risk Factors

Although risk factors for childhood and adolescent depression can be categorized as biologic, psychologic, or environmental (*Table 1*),

these factors are often intertwined.^{7,8,11,12} For example, parental depression is strongly associated with childhood and adolescent depression; children of parents with depression have a threefold greater risk of developing depression than those whose parents have no such history.^{13,14} Furthermore, the age when risk factors occur may predict future depression. Children diagnosed with a health condition such as diabetes mellitus or asthma between the ages of three and five years are likely to have a major depressive episode.¹² Likewise, children five years of age who were rated by teachers as being hostile were at greater risk of depression.¹²

Screening

The U.S. Preventive Services Task Force concluded that screening adolescents 12 to 18 years of age may lead to earlier detection and treatment of depression.¹⁵ There was insufficient evidence to recommend routine screening for younger children in primary care settings. The use of screening tools in children with at least one risk factor may be more helpful than universal screening. These tools can also indicate the severity of depressive symptoms. Screening tools include the Beck Depression Inventory for Primary

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>	<i>Comments</i>
Cognitive behavior therapy and interpersonal therapy should be used for the treatment of mild depression. Psychotherapy should be used in combination with medication for the treatment of moderate to severe depression in children and adolescents.	A	36, 39, 40	Meta-analysis, ³⁶ randomized controlled trials ^{39,40}
Tricyclic antidepressants should not be used in the treatment of childhood and adolescent depression.	A	46	Cochrane review
Fluoxetine (Prozac), citalopram (Celexa), and sertraline (Zoloft) are recommended as first-line treatments for childhood and adolescent depression.	C	47, 49, 50	Consensus guidelines
Treatment of major depression in children and adolescents should continue for at least six months.	C	51	Consensus guideline

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort.xml>.

Care for adolescents 12 to 18 years of age, and the Children's Depression Inventory for children and adolescents seven to 17 years of age.^{16,17} The Beck Depression Inventory for Primary Care is a self-report tool that includes seven questions that are scored from zero to three points each (Figure 1).^{17,18} A score of at least 4 warrants further evaluation for depression.¹⁹ The Children's Depression Inventory is available in self-report, parent-report, and teacher-report versions; scores are converted to a T-score, with scores of 65 or more indicating clinical significance.²⁰ Additional screening tools are described in Table 2.²¹⁻²⁴ Screening tools should be selected based on patient's age, reading level, and time available to complete the measures.

Presentation

Although diagnostic criteria for depression are the same for children and adults (Table 3²⁵), the manner in which these symptoms present may be different.²⁵ Adolescents

with depression are more likely to experience anhedonia, boredom, hopelessness, hypersomnia, weight change (including failure to reach appropriate weight milestones), alcohol or drug use, and suicide attempts. Younger children are more likely to have somatic symptoms, restlessness, separation anxiety, phobias, and hallucinations.²⁶ The child's cognitive level should be considered; for example, younger children may appear sad but have difficulty verbalizing their mood.²⁷ Differences have been found between parent reports and self-reports of depressive symptoms. Parents are more likely to indicate externalized symptoms such as irritability, whereas children are more likely to report internalized symptoms such as depressed mood.⁸

Diagnosis

When considering a diagnosis of depression, physicians must also consider likely medical causes of the presenting

Table 1. Risk Factors for Depressive Disorders in Children and Adolescents

Biologic	Psychological	Environmental
Family history of depression ^{7,8}	Emotional dependence ¹²	Antisocial peer group ¹¹
Female sex ¹¹	History of suicide attempts ^{11,12}	Decreased physical activity ¹¹
Hormonal changes during puberty ¹¹	Ineffective coping skills ^{11,12}	Increased parental conflict ^{8,12}
Low birth weight ⁷	Low self-esteem ^{11,12}	Loss of relationship (e.g., death of family member or friend, romantic relationship, friendship) ^{8,12}
Maternal age younger than 18 years ⁷	Negative body image ¹²	Low socioeconomic status ^{7,8}
Medical illness (e.g., asthma, diabetes mellitus, migraines) ⁷	Negative thinking styles (e.g., "Things like this always happen to me," "Nothing will ever go as planned") ^{11,12}	Overeating ¹¹
Obesity ¹¹	Self-consciousness ¹²	Poor academic performance ^{11,12}
Other psychological disorders (e.g., anxiety, learning disorders) ^{11,12}		Poor peer relationships ^{8,12}
Sleep disruptions ¹¹		Substance use ¹¹
		Traumatic event (e.g., physical or sexual abuse, accident) ⁸

Information from references 7, 8, 11, and 12.

Childhood and Adolescent Depression

symptoms, such as hyper- or hypothyroidism, anemia, or use of certain medications, including isotretinoin (Table 4).⁸ If the patient's mood is better explained by medical causes, the diagnosis of major depressive disorder is not appropriate.²⁵

Other psychological illnesses have presentations in children and adolescents that are similar to depression, especially adjustment, dysthymic, and bipolar disorders. Adjustment disorders are characterized by “a psychological response to an identifiable stressor or stressors.”²⁵ Symptoms develop within three

Children of parents with depression are three times more likely to develop depression.

months of the stressor and are in excess of what would be expected, or they cause significant impairment. Adjustment disorder with depressed mood can be mistaken for major depressive disorder; symptoms of both conditions include depressed mood, tearfulness, and feelings of hopelessness. However, the symptoms of adjustment disorder are related to a specific event and do not meet all criteria of a major depressive episode.²⁵

Dysthymic disorder in children and adolescents is characterized by depressed mood on most days for one year. Patients with dysthymia meet a subset of criteria for major depression, but the course of the illness is longer and the symptoms are less severe. Depressive symptoms may manifest as irritability in children and adolescents, and patients often have low self-esteem and poor social skills.

Bipolar disorder consists of periods of major depression alternating with mania or hypomania, during which patients have increased irritability, energy, grandiosity, and/or racing thoughts. It is important to rule out bipolar disorder when diagnosing depression, because antidepressant medications can initiate manic symptoms.

Risk Assessment for Suicide

The presence of a psychiatric disorder is one of the strongest correlates for suicide across age groups.²⁸ According to the Centers for Disease Control and Prevention, the incidence of suicide in children 10 to 14 years of age was approximately 0.9 per 100,000 in 2007, and the incidence in adolescents 15 to 19 years of age was 6.9 per 100,000.²⁹ The methods by which suicide was completed include firearms, suffocation, and poisoning; children most often use suffocation, whereas adolescents are more likely to use firearms.²⁹ Therefore, routine and ongoing assessment of suicidality in children and adolescents with depression is recommended, especially in patients who are receiving antidepressant medication.^{30,31}

Patient Self-Evaluation

Name: _____

Date: _____

This questionnaire consists of seven groups of statements. Read each group carefully, then circle the number next to the statement that best describes how you have felt during the past two weeks, including today. If several statements in the same group seem to apply equally well, choose the one with the highest number next to it.

1. Sadness

- 0 I do not feel sad.
- 1 I feel sad much of the time.
- 2 I am sad all the time.
- 3 I am so sad or unhappy I can't stand it.

2. Pessimism

- 0 I am not discouraged about my future.
- 1 I am more discouraged about my future than I used to be.
- 2 I do not expect things to work out for me.
- 3 I feel that my future is hopeless and will only get worse.

3. Past failure

- 0 I do not feel like a failure.
- 1 I have failed more than I should have.
- 2 As I look back, I see a lot of failure.
- 3 I feel that I am a total failure as a person.

4. Self-dislike

- 0 I feel the same about myself as I always have.
- 1 I have lost confidence in myself.
- 2 I am disappointed in myself.
- 3 I dislike myself.

5. Self-criticalness

- 0 I don't criticize or blame myself more than usual.
- 1 I am more critical of myself than I used to be.
- 2 I criticize myself for all of my faults.
- 3 I blame myself for everything bad that happens.

6. Suicidal thoughts or wishes

- 0 I don't have any thoughts of killing myself.
- 1 I have thoughts of killing myself, but I would not carry them out.
- 2 I would like to kill myself.
- 3 I would kill myself if I had the chance.

7. Loss of interest

- 0 I have not lost interest in other people or activities.
- 1 I am less interested in other people or things than before.
- 2 I have lost most of my interest in other people or things.
- 3 It's hard to get interested in anything.

Total score: _____

NOTE: 0 to 3 points = minimal symptoms of depression; 4 to 6 points = mild symptoms; 7 to 9 = moderate symptoms; 10 to 21 = severe symptoms. A high score alone does not indicate that a patient has a depressive disorder, but it does indicate that a more detailed evaluation should be performed. Physicians should review item 6 in particular, because it concerns suicidal ideation.¹⁸

Figure 1. Beck Depression Inventory for Primary Care.

Information from references 17 and 18.

Asking about suicide or using depression screening tools can be a means for assessing suicidal ideation in children and adolescents in the primary care setting. However, further inquiry into patient symptoms may be required because some tools lack specificity or are prone

Table 2. Screening Instruments for Childhood and Adolescent Depression

<i>Instrument</i>	<i>Age appropriateness (approximate years)</i>	<i>Reading level (grade)</i>	<i>Spanish version available?</i>	<i>Number of items</i>	<i>Time to complete (approximate minutes)</i>	<i>Sensitivity (%)</i>	<i>Specificity (%)</i>
Beck Depression Inventory	14 and older	6	Yes	21	5 to 10	84	81
Center for Epidemiological Studies Depression Scale	14 and older	6	No	20	5 to 10	84	75
Center for Epidemiological Studies Depression Scale for Children	12 to 18	6	Yes	20	5 to 10	71	57
Children's Depression Inventory	7 to 17	1	Yes	27	10 to 15	88	90
Patient Health Questionnaire-9	13 to 17	8	Yes	9	5 to 10	88	88
Reynolds Adolescent Depression Scale	13 to 18	3	No	30	10 to 15	89	90
Reynolds Child Depression Scale	8 to 12	2	Yes	30	10 to 15	97	73

Information from references 21 through 24.

Table 3. Criteria for Major Depressive Episode in Adults, Children, and Adolescents

The rights holder did not grant the American Academy of Family Physicians the right to sublicense this material to a third party. For the missing item, see the original print version of this publication.

Reprinted with permission from American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 4th ed., text rev. Copyright © 2000. Washington, DC: American Psychiatric Association; 2000:356.

to oversensitivity. If a patient has suicidal thoughts, further assessment should include questions about the frequency of thoughts; the presence and specificity of a plan; the lethality or availability of means to follow through with the plan; whether there are protective factors, such

as social support; and whether there are any other factors, such as substance use or a previous suicide attempt. If it is determined that a patient is at risk of self-harm, urgent referral to a mental health professional or emergency department is warranted.

Table 4. Differential Diagnosis of Depression

The rights holder did not grant the American Academy of Family Physicians the right to sublicense this material to a third party. For the missing item, see the original print version of this publication.

Adapted with permission from Richardson LP, Katzenellenbogen R. Childhood and adolescent depression: the role of primary care providers in diagnosis and treatment. Curr Probl Pediatr Adolesc Health Care. 2005;35(1):9.

Interventions for patients at lower risk of self-harm include involving parents or caregivers to provide close observation of the patient, and removing any weapons or means to self-harm. The frequency of follow-up visits may be increased, and patients may benefit from education on how to seek assistance after hours (e.g., emergency departments, crisis hotlines).^{8,30-32} Parents and family members should be educated about behaviors that would warrant immediate follow-up, such as increased suicidal ideation, impulsivity, irritability, restlessness, pressured speech, or psychomotor agitation.^{31,33}

Treatment

Treatment of childhood and adolescent depression consists of psychotherapy, pharmacotherapy, or a combination of these. Treatment should correspond to the level of depression, patient preferences, the developmental level of the patient, associated risk factors, and availability of services.⁸ Patient and family education about the associated risks and benefits of treatment, expectations regarding patient monitoring, and follow-up should be included.³³

PSYCHOTHERAPY

The American Psychiatric Association and the American Academy of Child and Adolescent Psychiatry recommend that psychotherapy always be a component of treatment for childhood and adolescent depression.³³ They recommend psychotherapy as an acceptable treatment option for patients with milder depression, and a combination of medication and psychotherapy in those with moderate to severe depression.

Cognitive behavior therapy (CBT) and interpersonal therapy have been proven effective in the treatment of adolescent depression, and CBT has been proven effective in the treatment of childhood depression. CBT usually consists of behavioral activation techniques and methods to increase coping skills, improve communication skills and peer relationships, solve problems, combat negative thinking patterns, and regulate emotions.^{8,34-36} In contrast, interpersonal therapy generally focuses on adapting to changes in relationships, transitioning personal roles, and forming interpersonal relationships.^{8,36} The effects of CBT on depressive symptoms are moderate,^{37,38} but it has not been proven more effective than placebo for treating acute

depression in adolescents.³⁹ A combination of CBT and medication has been shown to be more effective than medication alone in attaining remission of depression.^{37,40} Interpersonal therapy has not been compared with medication, combination treatment, or placebo, but it has been proven more effective than wait-list control groups with no therapy, and as effective or more effective than CBT.^{41,42}

PHARMACOTHERAPY

Although psychotherapy is a major component in the treatment of childhood and adolescent depression, adjunct use of medication is sometimes appropriate. Considerations for initiating antidepressant therapy include depression severity and history (Table 5).^{43,44} Extrapolation of adult data on antidepressant medications to children and adolescents may not be accurate, because neural pathways may not be fully developed, and serotonin and norepinephrine systems have different maturation rates.⁴⁵

Tricyclic antidepressants were previously used to treat depression in children, but studies have shown little to no benefit in adolescents and children.⁴⁶ Consensus guidelines recommend fluoxetine (Prozac), citalopram (Celexa), and sertraline (Zoloft) as first-line treatments for moderate to severe depression in children and adolescents.⁴⁷ A Cochrane review found that fluoxetine was the only agent with consistent evidence (from three randomized trials) that it is effective in

Table 5. Questions to Guide Initiation of Pharmacotherapy in Children and Adolescents with Depression

Is the depression of moderate to severe severity?
Has there been a prior episode of depression?
Has the patient been treated for depression with medication in the past?
Is there a family history of depression?
Is there a family history of depression with significant response to medication?
Have environmental stressors been modified with no associated improvements in mood?
Has evidence-based psychotherapy (i.e., cognitive behavior therapy, interpersonal therapy) been attempted without success?

NOTE: If the patient answers "yes" to any question, consider initiating pharmacotherapy.

Information from references 43 and 44.

decreasing depressive symptoms.⁴⁸ Escitalopram (Lexapro) is also licensed for the treatment of depression in adolescents 12 years and older. Treatment should begin at the lowest dosage available and titrated according to the patient's response and adverse effects. If initial first-line therapy is ineffective, another first-line agent should be considered. All antidepressants have a boxed warning for an increased risk of suicide; therefore, close monitoring is recommended (e.g., weekly telephone calls, scheduled visits for the first month of therapy) to assess for suicidality and other adverse effects, such as gastrointestinal effects, nervousness, headache, and restlessness.^{47,49} If there is limited improvement or no remission of symptoms after all first-line medications have been attempted, a psychiatric consultation is strongly recommended.⁵⁰

Treatment Duration and Referral

Treatment of depression in children and adolescents should continue for six months after remission.^{49,51} A double-blind, placebo-controlled trial of adolescents receiving fluoxetine found that those who received placebo after treatment had a shorter time to relapse than those who continued therapy.⁵² Patients in the fluoxetine group were significantly less likely to have a relapse of depressive symptoms (34 versus 60 percent). The decision to continue treatment for four to six months after remission of symptoms should be based on prior recurrent episodes of depression and current psychological and social stressors. Children younger than 11 years and those with chronic depression, comorbid substance use, psychiatric disorders, suicidality with plan, or lack of parental engagement in treatment should be referred to a psychiatrist.^{44,52}

Data Sources: We searched Essential Evidence Plus, the Agency for Healthcare Research and Quality evidence reports, the Cochrane Database of Systematic Reviews, Clinical Evidence, the National Guideline Clearinghouse, the U.S. Preventive Services Task Force, PubMed, Ovid Medline, and PsycINFO using the keywords childhood and adolescent depression, major depression and adolescents, major depression and childhood, major depression and children, depression and children, and depression and adolescents. The search included reviews, meta-analyses, randomized controlled trials, and clinical trials. Search dates: March 5 and 11, 2011.

The Authors

MOLLY S. CLARK, PhD, is an assistant professor of family medicine and director of the Health Psychology Fellowship at the University of Mississippi Medical Center, Jackson.

KATE L. JANSEN, PhD, is a health psychology fellow at the University of Mississippi Medical Center.

J. ANTHONY CLOY, MD, is an assistant professor of family medicine at the University of Mississippi Medical Center.

Address correspondence to Molly S. Clark, PhD, University of Mississippi Medical Center, 884 Lakeland Dr., Jackson, MS 39216 (e-mail: mclark@umc.edu). Reprints are not available from the authors.

Author disclosure: No relevant financial affiliations to disclose.

REFERENCES

1. Jane Costello E, Erkanli A, Angold A. Is there an epidemic of child or adolescent depression? *J Child Psychol Psychiatry*. 2006;47(12):1263-1271.
2. Weissman MM, Wolk S, Goldstein RB, et al. Depressed adolescents grown up. *JAMA*. 1999;281(18):1707-1713.
3. Brown J, Cohen P, Johnson JG, Smailes EM. Childhood abuse and neglect: specificity of effects on adolescent and young adult depression and suicidality. *J Am Acad Child Adolesc Psychiatry*. 1999;38(12):1490-1496.
4. Fergusson DM, Woodward LJ. Mental health, educational, and social role outcomes of adolescents with depression. *Arch Gen Psychiatry*. 2002;59(3):225-231.
5. Keenan-Miller D, Hammen CL, Brennan PA. Health outcomes related to early adolescent depression. *J Adolesc Health*. 2007;41(3):256-262.
6. Wells KB, Kataoka SH, Asarnow JR. Affective disorders in children and adolescents: addressing unmet need in primary care settings. *Biol Psychiatry*. 2001;49(12):1111-1120.
7. Kessler RC, Avenevoli S, Ries Merikangas K. Mood disorders in children and adolescents: an epidemiologic perspective. *Biol Psychiatry*. 2001;49(12):1002-1014.
8. Richardson LP, Katzenellenbogen R. Childhood and adolescent depression: the role of primary care providers in diagnosis and treatment. *Curr Probl Pediatr Adolesc Health Care*. 2005;35(1):6-24.
9. Cheung AH, Dewa CS, Levitt AJ, Zuckerbrot RA. Pediatric depressive disorders: management priorities in primary care. *Curr Opin Pediatr*. 2008;20(5):551-559.
10. Lieberman JA III. History of the use of antidepressants in primary care. *Primary Care Companion J Clin Psychiatry*. 2003;5(suppl 7):6-10.
11. Brent DA, Maalouf FT. Pediatric depression: is there evidence to improve evidence-based treatments? *J Child Psychol Psychiatry*. 2009;50(1-2):143-152.
12. Beardslee WR, Gladstone TR. Prevention of childhood depression: recent findings and future prospects. *Biol Psychiatry*. 2001;49(12):1101-1110.
13. Birmaher B, Ryan ND, Williamson DE, et al. Childhood and adolescent depression: a review of the past 10 years. Part I. *J Am Acad Child Adolesc Psychiatry*. 1996;35(11):1427-1439.

Childhood and Adolescent Depression

14. Birmaher B, Ryan ND, Williamson DE, Brent DA, Kaufman J. Childhood and adolescent depression: a review of the past 10 years. Part II. *J Am Acad Child Adolesc Psychiatry*. 1996;35(12):1575-1583.
15. U.S. Preventive Services Task Force. Screening and treatment for major depressive disorder in children and adolescents: recommendation statement [published correction appears in *Pediatrics*. 2009;123(6):1611]. *Pediatrics*. 2009;123(4):1223-1228.
16. Beck AT, Guth D, Steer RA, Ball R. Screening for major depression disorders in medical inpatients with the Beck Depression Inventory for Primary Care. *Behav Res Ther*. 1997;35(8):785-791.
17. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry*. 1961;4:561-571.
18. Kovacs M. *Children's Depression Inventory Manual*. North Tonawanda, NY: Multi-Health Systems, Inc.; 1992.
19. Winter LB, Steer RA, Jones-Hicks L, Beck AT. Screening for major depression disorders in adolescent medical outpatients with the Beck Depression Inventory for Primary Care. *J Adolesc Health*. 1999;24(6):389-394.
20. Timbremont B, Braet C, Driessens L. Assessing depression in youth: relation between the Children's Depression Inventory and a structured interview. *J Clin Child Adolesc Psychol*. 2004;33(1):149-157.
21. Sharp LK, Lipsky MS. Screening for depression across the lifespan: a review of measures for use in primary care settings. *Am Fam Physician*. 2002;66(6):1001-1008.
22. Richardson LP, McCauley E, Grossman DC, et al. Evaluation of the Patient Health Questionnaire-9 Item for detecting major depression among adolescents. *Pediatrics*. 2010;126(6):1117-1123.
23. Pignone M, Gaynes BN, Rushton JL, et al. Screening for depression: systematic evidence review. <http://www.ahrq.gov/downloads/pub/prevent/pdfser/depser.pdf>. Accessed August 3, 2011.
24. Reynolds WM, Mazza JJ. Reliability and validity of the Reynolds Adolescent Depression Scale with young adolescents. *J Sch Psychol*. 1998;36(3):295-312.
25. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed., text rev. Washington, DC: American Psychiatric Association; 2000.
26. Williams SB, O'Connor EA, Eder M, Whitlock EP. Screening for child and adolescent depression in primary care settings: a systematic evidence review for the US Preventive Services Task Force. *Pediatrics*. 2009;123(4):e716-e735.
27. Klein DN, Dougherty LR, Olino TM. Toward guidelines for evidence-based assessment of depression in children and adolescents. *J Clin Child Adolesc Psychol*. 2005;34(3):412-432.
28. Mościcki EK. Epidemiology of completed and attempted suicide: toward a framework for prevention. *Clin Neurosci Res*. 2001;1(5):310-323.
29. National Institute of Mental Health. Suicide in the U.S.: statistics and prevention. <http://www.mentalhealth.gov/health/publications/suicide-in-the-us-statistics-and-prevention/index.shtml>. Accessed March 11, 2011.
30. Horowitz LM, Ballard ED, Pao M. Suicide screening in schools, primary care and emergency departments. *Curr Opin Pediatr*. 2009;21(5):620-627.
31. Shain BN; American Academy of Pediatrics Committee on Adolescence. Suicide and suicide attempts in adolescents. *Pediatrics*. 2007;120(3):669-676.
32. Zuckerbrot RA, Cheung AH, Jensen PS, Stein RE, Laraque D; GLAD-PC Steering Group. Guidelines for adolescent depression in primary care (GLAD-PC): I. Identification, assessment, and initial management. *Pediatrics*. 2007;120(5):e1299-e1312.
33. American Psychiatric Association, American Academy of Child and Adolescent Psychiatry. The use of medication in treating childhood and adolescent depression: information for patients and families. <http://www.parentsmedguide.org/parentsmedguide.pdf>. Accessed March 5, 2011.
34. Fu CH, Williams SC, Cleare AJ, et al. Neural responses to sad facial expressions in major depression following cognitive behavioral therapy. *Biol Psychiatry*. 2008;64(6):505-512.
35. Weersing VR, Brent DA. Cognitive behavioral therapy for depression in youth. *Child Adolesc Psychiatr Clin N Am*. 2006;15(4):939-957.
36. David-Ferdon C, Kaslow NJ. Evidence-based psychosocial treatments for child and adolescent depression. *J Clin Child Adolesc Psychol*. 2008;37(1):62-104.
37. Brent D, Emslie G, Clarke G, et al. Switching to another SSRI or to venlafaxine with or without cognitive behavioral therapy for adolescents with SSRI-resistant depression: the TORDIA randomized controlled trial. *JAMA*. 2008;299(8):901-913.
38. Weisz JR, McCarty CA, Valeri SM. Effects of psychotherapy for depression in children and adolescents: a meta-analysis. *Psychol Bull*. 2006;132(1):132-149.
39. March J, Silva S, Petrycky S, et al. Fluoxetine, cognitive-behavioral therapy, and their combination for adolescents with depression: Treatment for Adolescents With Depression Study (TADS) randomized controlled trial. *JAMA*. 2004;292(7):807-820.
40. Kennard B, Silva S, Vitiello B, et al. Remission and residual symptoms after short-term treatment in the Treatment of Adolescents with Depression Study (TADS). *J Am Acad Child Adolesc Psychiatry*. 2006;45(12):1404-1411.
41. Klomek AB, Mufson L. Interpersonal psychotherapy for depressed adolescents. *Child Adolesc Psychiatr Clin N Am*. 2006;15(4):959-975.
42. Rosselló J, Bernal G. The efficacy of cognitive-behavioral and interpersonal treatments for depression in Puerto Rican adolescents. *J Consult Clin Psychol*. 1999;67(5):734-745.
43. Ryan ND. Child and adolescent depression: short-term treatment effectiveness and long-term opportunities. *Int J Methods Psychiatr Res*. 2003;12(1):44-53.
44. Emslie GJ, Mayes TL, Ruber M. Continuation and maintenance therapy of early-onset major depressive disorder. *Paediatr Drugs*. 2005;7(4):203-217.
45. Bylund DB, Reul AL. Childhood and adolescent depression: why do children and adults respond differently to antidepressant drugs? *Neurochem Int*. 2007;51(5):246-253.
46. Hazell P, O'Connell D, Heathcote D, Henry D. Tricyclic drugs for depression in children and adolescents. *Cochrane Database Syst Rev*. 2002;(2):CD002317.
47. Hughes CW, Emslie GJ, Crismon ML, et al. Texas Children's Medication Algorithm Project: update from Texas Consensus Conference Panel on Medication Treatment of Childhood Major Depressive Disorder. *J Am Acad Child Adolesc Psychiatry*. 2007;46(6):667-686.
48. Hetrick S, Merry S, McKenzie J, Sindahl P, Proctor M. Selective serotonin reuptake inhibitors (SSRIs) for depressive disorders in children and adolescents. *Cochrane Database Syst Rev*. 2007;(3):CD004851.
49. Cincinnati Children's Hospital Medical Center. Best evidence statement: treatment of children and adolescents with major depressive disorder (MDD) during the acute phase. <http://www.cincinnatichildrens.org/service/janderson-center/evidence-based-care/bests>. Accessed November 28, 2011.
50. Cheung AH, Zuckerbrot RA, Jensen PS, Ghalib K, Laraque D, Stein RE; GLAD-PC Steering Group. Guidelines for adolescent depression in primary care (GLAD-PC): II. Treatment and ongoing management. *Pediatrics*. 2007;120(5):e1313-e1326.
51. Emslie GJ, Mayes TL. Mood disorders in children and adolescents: psychopharmacological treatment. *Biol Psychiatry*. 2001;49(12):1082-1090.
52. Emslie GJ, Heiligenstein JH, Hoog SL, et al. Fluoxetine treatment for prevention of relapse of depression in children and adolescents: a double-blind, placebo-controlled study. *J Am Acad Child Adolesc Psychiatry*. 2004;43(11):1397-1405.

Copyright of American Family Physician is the property of American Academy of Family Physicians and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.